

## Details




Molecular Ecology  
Early View

## ARTICLE

Barriers to chimpanzee gene flow at the south-east edge of their distribution

[View article page](#)

Noémie Bonnin, Alex K. Piel, Richard P. Brown, Yingying Li, Andrew Jesse Connell, ... [See all authors](#)



CITE



Check for updates

© 2023 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd.

<https://doi.org/10.1111/mec.16986>

ISSN 0962-1083

eISSN 1365-294X

Accepted 2 May 2023

Revised 28 April 2023

Received 12 September 2022

Pages n/a - n/a

## Abstract

Populations on the edge of a species' distribution may represent an important source of adaptive diversity, yet these populations tend to be more fragmented and are more likely to be geographically isolated. Lack of genetic exchanges between such populations, due to barriers to

Received: 12 September 2022 | Revised: 28 April 2023 | Accepted: 2 May 2023

DOI: 10.1111/mec.16986

## ORIGINAL ARTICLE

MOLECULAR ECOLOGY WILEY

## Barriers to chimpanzee gene flow at the south-east edge of their distribution

Noémie Bonnin<sup>1</sup> | Alex K. Piel<sup>2</sup> | Richard P. Brown<sup>1</sup> | Yingying Li<sup>3</sup> | Andrew Jesse Connell<sup>3</sup> | Alexa N. Avitto<sup>3</sup> | Jean P. Boubli<sup>4</sup> | Adrienne Chitayat<sup>5</sup> | Jasmin Giles<sup>3</sup> | Madhurima S. Gundlapally<sup>3</sup> | Iddi Lipende<sup>6</sup> | Elizabeth V. Lonsdorf<sup>7,8</sup> | Deus Mjungu<sup>9</sup> | Dismas Mwacha<sup>9</sup> | Lilian Pintea<sup>10</sup> | Anne E. Pusey<sup>11</sup> | Jane Raphael<sup>12</sup> | Serge A. Wich<sup>1,5</sup> | Michael L. Wilson<sup>13,14,15</sup> | Emily E. Wroblewski<sup>16</sup> | Beatrice H. Hahn<sup>3</sup> | Fiona A. Stewart<sup>1,2</sup>

<sup>1</sup>School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK

<sup>2</sup>Department of Anthropology, University College London, London, UK

<sup>3</sup>Departments of Medicine and Microbiology, University of Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>4</sup>School of Science, Engineering & Environment, University of Salford, Salford, UK

<sup>5</sup>Institute of Biodiversity and Ecological Dynamics, University of Amsterdam, Amsterdam, The Netherlands

<sup>6</sup>Tanzania Wildlife Research Institute (TAWIRI), Arusha, Tanzania

<sup>7</sup>Department of Psychology, Franklin and Marshall College, Lancaster, Pennsylvania, USA

<sup>8</sup>Department of Anthropology, Emory University, Atlanta, Georgia, USA

<sup>9</sup>Gombe Stream Research Centre, The Jane Goodall Institute–Tanzania, Kigoma, Tanzania

<sup>10</sup>Conservation Science Department, The Jane Goodall Institute, Washington, District of Columbia, USA

<sup>11</sup>Department of Evolutionary Anthropology, Duke University, Durham, North Carolina, USA

<sup>12</sup>Gombe National Park, Kigoma, Tanzania

<sup>13</sup>Department of Anthropology, University of Minnesota, Minneapolis, Minnesota, USA

<sup>14</sup>Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota, USA

<sup>15</sup>Institute on the Environment, University of Minnesota, St. Paul, Minnesota, USA

<sup>16</sup>Anthropology Department, Washington University, St. Louis, Missouri, USA

## Correspondence

Noémie Bonnin, School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK.  
Email: [noemie.bonnin@gmail.com](mailto:noemie.bonnin@gmail.com)

## Abstract

Populations on the edge of a species' distribution may represent an important source of adaptive diversity, yet these populations tend to be more fragmented and are more